



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

The experience-dependent increase in deep sleep activity is reduced in children with attention-deficit/hyperactivity disorder

Furrer, Melanie ; Ringli, Maya ; Kurth, Salome ; Brandeis, Daniel ; Jenni, Oskar G ; Huber, Reto

Abstract: Objective/Background Learning of a visuomotor adaptation task during wakefulness leads to a local increase in slow-wave activity (SWA, EEG power between 1 and 4.5 Hz) during subsequent deep sleep. Here, we examined this relationship between learning and SWA in children with attention-deficit/hyperactivity disorder (ADHD). Patients/Methods Participants were 15 children with ADHD (9.7 – 14.8 y, one female) and 15 age-matched healthy controls (9.6 – 15.7 y, three female). After the completion of a visuomotor adaptation task in the evening, participants underwent an all-night high-density (HD, 128 electrodes) sleep-EEG measurement. Results Healthy control children showed the expected right-parietal increase in sleep SWA after visuomotor learning. Despite no difference in visuomotor learning, the local up-regulation during sleep was significantly reduced in ADHD patients compared to healthy controls. Conclusions Our results indicate that the local, experience-dependent regulation of SWA is different in ADHD patients. Because the customarily observed heightened regulation in children was related to sensitive period maturation, ADHD patients may lack certain sensitive periods or show a developmental delay.

DOI: <https://doi.org/10.1016/j.sleep.2019.09.018>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-177710>

Journal Article

Accepted Version

Originally published at:

Furrer, Melanie; Ringli, Maya; Kurth, Salome; Brandeis, Daniel; Jenni, Oskar G; Huber, Reto (2020).

The experience-dependent increase in deep sleep activity is reduced in children with attention-deficit/hyperactivity disorder. *Sleep Medicine*, 75:50-53.

DOI: <https://doi.org/10.1016/j.sleep.2019.09.018>

Journal Pre-proof

The experience-dependent increase in deep sleep activity is reduced in children with attention-deficit/hyperactivity disorder

Melanie Furrer, Maya Ringli, Salome Kurth, Daniel Brandeis, Oskar G. Jenni, Reto Huber



PII: S1389-9457(19)30331-4

DOI: <https://doi.org/10.1016/j.sleep.2019.09.018>

Reference: SLEEP 4199

To appear in: *Sleep Medicine*

Received Date: 26 April 2019

Revised Date: 30 August 2019

Accepted Date: 1 September 2019

Please cite this article as: Furrer M, Ringli M, Kurth S, Brandeis D, Jenni OG, Huber R, The experience-dependent increase in deep sleep activity is reduced in children with attention-deficit/hyperactivity disorder, *Sleep Medicine*, <https://doi.org/10.1016/j.sleep.2019.09.018>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The experience-dependent increase in deep sleep activity is reduced in children with attention-deficit/hyperactivity disorder.

Melanie Furrer^{1,2}, Maya Ringli¹, Salome Kurth³, Daniel Brandeis^{4,5,6}, Oskar G. Jenni^{1,2,6}, Reto Huber^{1,2,4}

1. Child Development Center, University Children's Hospital Zurich, Steinwiesstrasse 75, 8032 Zurich, Switzerland
2. Children's Research Center, University Children's Hospital Zurich, Steinwiesstrasse 75, 8032 Zurich, Switzerland
3. Pulmonary Clinic, University Hospital Zurich, Rämistrasse 100, 8091 Zurich, Switzerland
4. Department of Child and Adolescent Psychiatry and Psychotherapy, Psychiatric Hospital, University of Zurich, Lenggstrasse 31, 8032 Zurich, Switzerland
5. Department of Child and Adolescent Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, J5, 68159 Mannheim, Germany
6. Center for Integrative Human Physiology, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

melanie.furrer@kispi.uzh.ch; maya.ringli-gruenenwald@insel.ch; salome.kurth@usz.ch;
brandeis@kjpd.uzh.ch; oskar.jenni@kispi.uzh.ch

Corresponding author:

Prof. Reto Huber, PhD

University Children's Hospital

Steinwiesstrasse 75

8032 Zürich

Switzerland

phone: +41 44 266 81 60

fax: +41 44 266 78 66

e-mail: reto.huber@kispi.uzh.ch

Abstract*Objective/Background*

Learning of a visuomotor adaptation task during wakefulness leads to a local increase in slow-wave activity (SWA, EEG power between 1 and 4.5 Hz) during subsequent deep sleep. Here, we examined this relationship between learning and SWA in children with attention-deficit/hyperactivity disorder (ADHD).

Patients/Methods

Participants were 15 children with ADHD (9.7 – 14.8 y, one female) and 15 age-matched healthy controls (9.6 – 15.7 y, three female). After the completion of a visuomotor adaptation task in the evening, participants underwent an all-night high-density (HD, 128 electrodes) sleep-EEG measurement.

Results

Healthy control children showed the expected right-parietal increase in sleep SWA after visuomotor learning. Despite no difference in visuomotor learning, the local up-regulation during sleep was significantly reduced in ADHD patients compared to healthy controls.

Conclusions

Our results indicate that the local, experience-dependent regulation of SWA is different in ADHD patients. Because the customarily observed heightened regulation in children was related to sensitive period maturation, ADHD patients may lack certain sensitive periods or show a developmental delay.

Keywords

ADHD; high-density EEG; slow-wave activity; development; sensitive period maturation; local sleep

1. Introduction

The need for sleep is best reflected in slow-wave activity (SWA, EEG power between 1 and 4.5 Hz) during deep sleep (Achermann and Borbély, 2017). There is compelling evidence indicating that the build-up of sleep need is dependent on daytime experience (Diekelmann and Born, 2010; Tononi and Cirelli, 2019). For example, SWA is up-regulated over the right-parietal cortex after learning a visuomotor adaptation task (Huber et al., 2004; Määttä et al., 2010), suggesting that sleep need (i.e., SWA) increases as a function of experience-dependent plasticity. This change in SWA supposedly reflects plastic changes underlying the consolidation of the learned task in the brain area crucial for visuomotor integration (Ghilardi et al., 2000). Note that deep sleep SWA is considerably higher in children than in adults (Campbell and Feinberg, 2009; Kurth et al., 2010). Correspondingly, during development, when the maturing brain shows the highest plasticity, the experience-dependent increase in SWA is most pronounced (Wilhelm et al., 2014). Together, these findings indicate that local experience-dependent regulation of SWA reflects sensitive periods during maturation.

Children with attention-deficit/hyperactivity disorder (ADHD) show an anterior-posterior imbalance as well as a global reduction in SWA compared to healthy age-matched children (Ringli et al., 2013; Furrer et al., submitted). Thus, we tested whether the local increase in SWA following a visuomotor adaptation task is less pronounced in ADHD compared to healthy participants, to provide insights into the role of deep sleep in ADHD.

2. Methods

1.1 Participants

To test this hypothesis, 15 children with ADHD (9.7 – 14.8 y, mean: 12.4 y, 1 female, 1 left-handed) meeting the DSM-IV criteria were recruited from the Department for Child and Adolescent Psychiatry of the University of Zurich (data of 9 subjects published previously in Ringli et al., (2013)). Eight ADHD patients received no psychostimulant medication during the time of their participation in the study. Seven of these children have never received psychostimulants, and one participant was treated with methylphenidate until two years before the start of the study. The remaining seven ADHD patients were receiving methylphenidate (Concerta, Ritalin) on a regular basis (mean: 45.29 mg/day, range: 15 – 80 mg/day). One of these patients refrained from the intake of medication on the day of the sleep assessment.

15 sex- and age-matched healthy controls (9.6 – 15.7 y, mean: 12.3 y, three female, all right-handed) were included from a study published previously by (Wilhelm et al., 2014). Healthy controls were never diagnosed with a psychiatric disorder and did not take any medication during the period of assessments.

Six months prior to the study, no participant traveled across more than one time zone. During the seven days before the sleep assessment, a regular sleep-wake rhythm was maintained as verified with wrist accelerometry (Actiwatch Plus, AW4, Cambridge Neurotechnology) and sleep diaries. Written informed consent was obtained from a parent, the study was approved by the local ethics committee and performed according to the Declaration of Helsinki.

1.2 Procedures

All participants underwent an all-night high spatial resolution (high-density, HD) sleep-EEG measurement (128 electrodes) in the sleep laboratory of the University Children's Hospital Zurich (preprocessing of EEG-data in Supplementary Methods). Healthy control participants underwent two night recordings (rotation and baseline condition) separated by at least one week with the order of conditions balanced across subjects. In the evening of each condition, they performed a visuomotor adaptation task in which they moved a cursor from a central starting point to one of four target points in a pseudorandomized order. In the rotation condition, the visual feedback of the target points was rotated at a fixed angle. The learning session consisted of four rotation-blocks (15°, 30°, 45°, and 60°), each comprising three trials with the same rotation angle. In each of the trials, subjects had to execute 44 movements. After each trial, subjects received visual feedback about their performance. Learning was parametrized by measuring the directional error at peak velocity (Fig. 1a). In the baseline condition, participants performed the same number of movements without any rotation. A detailed description of the task can be found in (Wilhelm et al., 2014). As a major limitation of the study, ADHD patients only participated in the rotation condition.

1.3 Data analysis

SWA was averaged over the first hour of artifact-free NREM sleep because this yields a reliable estimate of deep sleep brain activity (Lustenberger et al., 2017). As in previous investigations addressing local sleep EEG topography (Ringli et al., 2013; Wilhelm et al., 2014), SWA was normalized by dividing the value of each electrode by the mean across all electrodes. To investigate the learning-induced increase in right-parietal SWA, the brain area crucial for visuomotor integration

(Ghilardi et al., 2000), during the rotation condition independent of a baseline recording, electrode-wise one-sided paired t-tests were applied to compare the right- with the left brain hemisphere. To disentangle possible naturally existing hemispheric asymmetries in the level of SWA, we performed this analysis for the baseline condition.

To account for multiple testing, we defined a cluster of electrodes as minimally six significant neighboring electrodes, because when calculating statistical tests across electrodes (one-sided, significance level of 5 %), five electrodes will become significant by chance. A similar approach was used previously (Wilhelm et al., 2014).

3. Results

We first analyzed the learning efficiency during the task before sleep and found that ADHD patients showed no differences in the learning curve compared to healthy controls. Furthermore, neither the rhythmicity nor straightness of the executed movements differed between the groups, indicating that the cognitive effort to complete this task was similar in ADHD and control participants (Fig. 1b, see also Supplementary Table S1, Supplementary Methods for more information). Thus, we would expect similar experience-dependent plastic changes in both groups before sleep.

Next, we confirmed previous findings of locally increased parietal SWA in the rotation- compared to the baseline condition (Wilhelm et al., 2014) during the first hour of artifact-free NREM sleep in the control group (Fig. 1c, top). To overcome the study design limitation in ADHD patients, we used the left hemisphere of the rotation condition as a "baseline," which is a way to estimate the experience-dependent SWA change without a baseline recording (Poryazova et al., 2015). When performing this comparison in healthy controls, we found a cluster of twelve electrodes with significantly increased SWA over the right cortex (Fig. 1c, bottom left). To exclude pre-existing asymmetries, we performed the same comparison for the baseline condition (Fig. 1c, bottom right). Visual inspection of this comparison suggests that the SWA increase over the parietal cortex is exclusive to the rotation condition. The positive correlation between the right-parietal SWA increase after the rotation compared to baseline and the local difference in SWA when comparing the right to the left hemisphere in the rotation condition only ($r = 0.78$, $p = 0.0006$) supports our approach.

Thus, we used this "right-left hemisphere" approach to test if ADHD patients differ from healthy controls in terms of the experience-dependent local SWA increase. Electrode-wise 2 x 2 ANOVA with the factor group and the within-group factor brain hemisphere revealed no significant main effect of

group or brain hemisphere, but a significant interaction between the two in a cluster of six parietal electrodes ($F(1,28) = 10.68$, $p = 0.0029$), indicating that ADHD patients differ from healthy controls in the experience-dependent increase of local SWA (Fig. 1d). Indeed, a direct comparison showed that the local change in SWA is reduced in ADHD patients compared to healthy controls (Fig. 1e). The small sample size limited the analyses of the effects of medication, gender, and handedness (see Supplementary Results). Sleep efficiency and architecture did not differ between groups, while total time in bed was longer in ADHD than control participants (Supplementary Table S2). Because only the first hour of NREM sleep was analyzed, this should not have influenced the results. Potential first night effects could not be assessed in the ADHD group because ADHD patients spent only one night in the sleep laboratory. In the control group, however, first and second nights did not differ in terms of sleep architecture nor SWA topography (data are not shown).




Figure 1. (a) Experimental design. Subjects performed a visuomotor adaptation task in the evening before an all-night sleep HD-EEG recording. Directional error represents the angle between the line from the initial hand position to the position of the target and the line to the position of the hand at the peak outward velocity. (b) Learning curves with normalized directional error (divided by rotation angle). ADHD patients ($n = 15$) and healthy controls ($n = 15$) improved across trials and rotation blocks ($p <$

0.0001 for main effect of trial and block; $p = 0.8078$ for main effect of group). (c) Top: T-values resulting from electrode-wise Student's one-tailed paired t-tests comparing SWA (first hour of artifact-free NREM sleep) of the baseline session with SWA of the rotation session in healthy controls. Bottom: T-values comparing SWA of the right hemisphere with SWA of the left hemisphere in healthy controls for the rotation and the baseline sessions. White dots represent electrodes with a significant SWA increase ($p < 0.05$). (d) Electrode-wise interaction between brain hemisphere and group. F-values are represented on the scalp map with significant electrodes in white. (e) Topographical distribution of the difference between right and left SWA and boxplots displaying the mean difference over the cluster of six electrodes showing a significant interaction.

4. Discussion

Learning a task during wakefulness causes a build-up of sleep need, which is best reflected in the initial increase of SWA (e.g., the first hour of sleep) during subsequent sleep. Even though ADHD patients learned a visuomotor adaptation task equally well as their normally developing peers, we observed no corresponding build-up of SWA. Hence, the general reduction in SWA found in ADHD patients (Furrer et al., submitted) could be the result of a reduced experience-dependent build-up of sleep need. As the heightened experience-dependent regulation of SWA in children was related to sensitive period maturation (Wilhelm et al., 2014), ADHD patients may lack certain sensitive periods or simply show a developmental delay.

Our results may also relate to behavioral consequences, since the up-regulation of SWA after learning is necessary for sleep-dependent performance gains (Huber et al., 2004; Landsness et al., 2009). Notably, the memory consolidating effects of sleep were not found in ADHD patients (Prehn-Kristensen et al., 2013, 2011). Moreover, boosting SWA by transcranial oscillatory direct current stimulation, presumably bringing it to normal levels, improved declarative memory consolidation in children with ADHD (Prehn-Kristensen et al., 2014). Unfortunately, we did not assess task performance after sleep in our participants and therefore, cannot draw any direct conclusions related to sleep-dependent performance.

1. Acknowledgments

This work was supported by the Swiss National Science Foundation (grant number PPOOA-114923), the Clinical Research Priority Program (CRPP) "Sleep and Health" of the University of Zurich Switzerland and the HMZ Flagship Project "SleepLoop" of University Medicine Zurich Switzerland.

The authors thank Soraya Souissi, Robert Aellen, and Katharina Böttger for their help in data collection and preprocessing, and Susanne Staubli for her support in graphical illustration.

2. References

- Achermann, P., Borbély, A.A., 2017. Sleep homeostasis and models of sleep regulation, in Kryger, M.H., Roth, T., Dement, W.C. (Eds.), *Principles and Practice of Sleep Medicine*. Philadelphia: Elsevier, pp. 377–387.
- Campbell, I.G., Feinberg, I., 2009. Longitudinal trajectories of non-rapid eye movement delta and theta EEG as indicators of adolescent brain maturation. *Proc Natl Acad Sci U S A* 106, 5177–5180. <https://doi.org/10.1073/pnas.0812947106>
- Diekelmann, S., Born, J., 2010. The memory function of sleep. *Nat. Rev. Neurosci.* 11, 114–126. <https://doi.org/10.1038/nrn2762>
- Ghilardi, M.-F., Ghez, C., Dhawan, V., Moeller, J., Mentis, M., Nakamura, T., Antonini, A., Eidelberg, D., 2000. Patterns of regional brain activation associated with different forms of motor learning. *Brain Res.* 871, 127–145. [https://doi.org/10.1016/S0006-8993\(00\)02365-9](https://doi.org/10.1016/S0006-8993(00)02365-9)
- Huber, R., Ghilardi, M.F., Massimini, M., Tononi, G., 2004. Local sleep and learning. *Nature* 430, 78–81. <https://doi.org/10.1038/nature02663>
- Kurth, S., Ringli, M., Geiger, A., LeBourgeois, M., Jenni, O.G., Huber, R., 2010. Mapping of cortical activity in the first two decades of life: a high-density sleep electroencephalogram study. *J Neurosci* 30, 13211–13219. <https://doi.org/10.1523/jneurosci.2532-10.2010>
- Landsness, E.C., Crupi, D., Hulse, B.K., Peterson, M.J., Huber, R., Ansari, H., Coen, M., Cirelli, C., Benca, R.M., Ghilardi, M.F., Tononi, G., 2009. Sleep-dependent improvement in visuomotor learning: a causal role for slow waves. *Sleep* 32, 1273–1284.
- Lustenberger, C., Mouthon, A.-L., Tesler, N., Kurth, S., Ringli, M., Buchmann, A., Jenni, O.G., Huber, R., 2017. Developmental trajectories of EEG sleep slow-wave activity as a marker for motor skill development during adolescence: a pilot study. *Dev. Psychobiol.* 59, 5–14. <https://doi.org/10.1002/dev.21446>
- Määttä, S., Landsness, E., Sarasso, S., Ferrarelli, F., Ferreri, F., Ghilardi, M.F., Tononi, G., 2010. The effects of morning training on night sleep: A behavioral and EEG study. *Brain Res. Bull.* 82, 118–123. <https://doi.org/10.1016/j.brainresbull.2010.01.006>
- Poryazova, R., Huber, R., Khatami, R., Werth, E., Brugger, P., Barath, K., Baumann, C.R., Bassetti, C.L., 2015. Topographic sleep EEG changes in the acute and chronic stage of hemispheric stroke. *J. Sleep Res.* 24, 54–65. <https://doi.org/10.1111/jsr.12208>
- Prehn-Kristensen, A., Göder, R., Fischer, J., Wilhelm, I., Seeck-Hirschner, M., Aldenhoff, J., Baving, L., 2011. Reduced sleep-associated consolidation of declarative memory in attention-deficit/hyperactivity disorder. *Sleep Med.* 12, 672–679. <https://doi.org/10.1016/J.SLEEP.2010.10.010>
- Prehn-Kristensen, A., Munz, M., Göder, R., Wilhelm, I., Korr, K., Vahl, W., Wiesner, C.D., Baving, L., 2014. Transcranial Oscillatory Direct Current Stimulation During Sleep Improves Declarative Memory Consolidation in Children With Attention-deficit/hyperactivity Disorder to a Level Comparable to Healthy Controls. *Brain Stimul.* 7, 793–799.

<https://doi.org/10.1016/J.BRS.2014.07.036>

Prehn-Kristensen, A., Munz, M., Molzow, I., Wilhelm, I., Wiesner, C.D., Baving, L., 2013. Sleep Promotes Consolidation of Emotional Memory in Healthy Children but Not in Children with Attention-Deficit Hyperactivity Disorder. *PLoS One* 8, e65098.

<https://doi.org/10.1371/journal.pone.0065098>

Ringli, M., Souissi, S., Kurth, S., Brandeis, D., Jenni, O.G., Huber, R., 2013. Topography of sleep slow-wave activity in children with attention-deficit/hyperactivity disorder. *Cortex* 49, 340–347.

<https://doi.org/10.1016/j.cortex.2012.07.007>

Tononi, G., Cirelli, C., 2019. Sleep and synaptic down-selection. *Eur. J. Neurosci.*

<https://doi.org/10.1111/ejn.14335>

Wilhelm, I., Kurth, S., Ringli, M., Mouthon, A.-L., Buchmann, A., Geiger, A., Jenni, O.G., Huber, R., 2014. Sleep slow-wave activity reveals developmental changes in experience-dependent plasticity. *J. Neurosci.* 34, 12568–75. <https://doi.org/10.1523/JNEUROSCI.0962-14.2014>

3. Declaration of interest

None.

4. Supplementary Material

Supplementary Table S1

Supplementary Table S2

Supplementary Results

Supplementary Methods

- EEG slow-wave activity (SWA) during sleep reflects experience-dependent plasticity.
- Visuomotor learning before sleep was normal in children with ADHD.
- The experience-dependent SWA increase during subsequent sleep was reduced in ADHD.